

TEAMWORK IMPROVES SAFETY

<u>Signatures</u>
1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17

KCA TOOL BOX TALK: Lead Exposure

Exposure to lead fumes and particles can come from many sources, including abrasive blasting, sanding, torch cutting, scraping, and loosening old paint with a propane torch. Dust and fumes can be inhaled, including by family members shaking out clothes. Lead can also get into the body by being transferred from dirty hands to food and drink. Serious damage can occur to your lungs, brain, liver and other organs. Children are particularly at risk for lead dust brought home from the job.

Here is an Example: Joe was removing lead paint on a highway bridge job. After a few months, he went to the hospital because he was experiencing difficulty concentrating, vomiting and weight loss. The doctor diagnosed high levels of lead in his blood.

- 1. What protective equipment should he have worn while working?
- 2. Have you ever experienced any of the symptoms mentioned while working in a lead exposure area?
- 3. If so, did you go to the doctor for a blood test to check your level of lead exposure?

Preventing Injuries from Lead Exposure:

- Never abrasive blast or torch cut painted metal structures until they are tested for lead based paint. OSHA has a standard for lead in construction work.
- Wash your hands and face before eating, drinking or smoking and never eat, or use tobacco products in the work area.
- Wear personal protective equipment when there is a lead exposure risk. You
 may need coveralls, gloves, shoe coverlets and vented goggles/face shield or
 respirator if high exposure.

For more information visit OSHA REGULATIONS: 1926.62

www.KeystoneContractors.com